

AD-A034 176

BOSTON UNIV MASS DEPT OF CHEMISTRY
PHOTOISOMERIZATION OF LINKED ANTHRACENSES. A DEMANDING TEST OF --ETC(U)
DEC 76 W R BERGMARK, G JONES

N00014-76-C-0442

F/G 7/3

NL

UNCLASSIFIED


1 OF 1
AD
A034176



END

DATE
FILMED

2-77



ADA034176

OFFICE OF NAVAL RESEARCH

Contract N00014-76-C-0442

Project No. NR 051-574

TECHNICAL REPORT NO. 5

PHOTOISOMERIZATION OF LINKED ANTHRACENES. A DEMANDING TEST
OF MOLECULAR GEOMETRY FOR INTERMEDIATES IN PHOTODIMERIZATION


by

W.R. Bergmark and G. Jones, II

Prepared for Publication
in the
Journal of the American Chemical Society

Department of Chemistry
Boston University
Boston, Massachusetts 02215

December 1, 1976



Reproduction in whole or in part is permitted for any purpose
of the United States Government

Approved for public release; distribution unlimited

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|---|---|---|
| 1. REPORT NUMBER (9) Technical Report No. 5, 1 NOV 75-31 Aug 76 | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) (6) Photoisomerization of Linked Anthracenes. A Demanding Test of Molecular Geometry for Intermediates in Photodimerization. | 5. TYPE OF REPORT & PERIOD COVERED Technical, 11/1/75-8/31/76 | |
| 6. PERFORMING ORG. REPORT NUMBER | | 8. CONTRACT OR GRANT NUMBER(s) (15) N00014-76-C-0442 new |
| 7. AUTHOR(s) (10) William R. Bergmark and G. Jones, II | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-574/9-26-75 (472) | |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry, Boston University Boston, Massachusetts 02215 | 12. REPORT DATE December 1, 1976 | |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 DODAAD Code N00014 | 13. NUMBER OF PAGES 8 (12) 11 p | |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | 15. SECURITY CLASS. (of this report) unclassified | |
| 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE | | |
| 16. DISTRIBUTION STATEMENT (of this Report) Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release; distribution unlimited. | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES Prepared for publication in the Journal of the American Chemical Society | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Anthracene photodimerization, bichromophoric molecules, excimers, sandwich dimers, diradicals, fluorescence lifetimes and quantum yields. | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Valence photoisomerization and photophysical properties of a series of dianthrylethanes, dianthrylmethanes, and tetrabenzo(2.2)paracyclophane are reported. Fluorescence lifetimes and quantum yields depend upon the nature of the linkage between anthracene moieties. Quantum efficiency for photoisomerization depends on the substitution pattern at 10,10 positions. Sandwich dimers with excimer-like interaction are generated on low temperature photolysis in glasses of dianthrylethane but not dianthrylmethane. It is concluded that geometrical requirements for excimerization and photodimerization are quite different. | | |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

401003

PHOTOISOMERIZATION OF LINKED ANTHRACENES. A DEMANDING TEST OF
MOLECULAR GEOMETRY FOR INTERMEDIATES IN PHOTODIMERIZATION

ABSTRACT

Valence photoisomerization and photophysical properties of dianthrylmethanes, (lc-e), dianthrylethanes (la,b), and tetrabenzo(2.2)paracyclophane (lf) have been examined. Fluorescence yields and lifetimes are significantly a function of the bridging group for the linked anthracenes [e.g., $\gamma_f(\text{ethanes}) > \gamma_f(\text{methanes})$]. Quantum yields for photoisomerization (ϕ_{12}), on the other hand, appear to be more a function of the 9' substituent group Y. Evidence for emission from a weakly stabilized excimer-like state is presented for la, whereas similar experiments on lc-e in which "sandwich dimers" are produced by 254 nm photolysis at 77 K reveal only normal anthracene fluorescence. The effects of structure on photoreactivity and singlet excited state decay for the linked anthracenes are mechanistically consistent with singlet lifetime-limiting formation of diradical intermediates which partition to reactant and photoisomer. The geometrical requirements for excimerization exceed those for photoreaction of proximal anthracenes, suggesting a subordinate role for excimers in photodimerization.

| | |
|----------------------------|---------------------------------------|
| ACCESSION NO. | |
| RTIS | NOT INDEXED <input type="checkbox"/> |
| CRC | DATE INDEXED <input type="checkbox"/> |
| UNANALYZED | |
| IDENTIFICATION | |
| BY | |
| DISTRIBUTION/ANALYSIS CODE | |
| DATE | FILE |
| A | |

Photoisomerization of Linked Anthracenes. A Demanding Test of Molecular
Geometry for Intermediates in Photodimerization.


Sir:

We have examined the photoisomerization and emission characteristics of linked anthracenes 1, which represent a graded series of molecular constriction ending with the rigid tetrabenzo[2.2]paracyclophane, 1f. Our results are informative with regard to the nature of necessary intermediates in anthracene photodimerization. The data are presented in Table I.



Excimers. Sandwich-like excimer states have long been assumed (by virtue of similarity in structure to photodimers) to be intermediates in anthracene photodimerizations, and a number of experiments have been interpreted as supporting this view.⁴ Examination of models of the linked systems 1, in particular those with a one-carbon bridge (1c-e), reveals that the intramolecular counterparts to excimers should be strained, beyond the normally expected^{4d} 5-10 kcal/mol excimer binding energy if they assume a sandwich-like geometry.

The synthesis of sandwich dimers is possible using the clever experimental approach of Chandross and Ferguson⁵ involving photolysis of anthracene photodimers at 77°K. We have applied this procedure to compounds 1c-e and find only normal fluorescence (that observed for 1c-e in glasses at 77°K) after irradiation (254 nm) of photoisomers 2c-e in 2-methyltetrahydrofuran glasses. Unusual or long wavelength excimer emission found for a number of sandwich dimers^{5,6} is not observed, even on using a more rigid hydrocarbon glass⁶ (3:2, methylcyclohexane/decalin).


Rather different results are provided by 1,2-bis(9-anthryl)ethane, 1a. After its photoisomer 2a is irradiated at 254 nm in a rigid glass at 77°K, a strong emission is observed, the major part of which is very similar to the normal 77°K emission spectrum except for an 8 nm red-shift. Melting and refreezing the glass restores the normal spectrum. We interpret the red-shifted emission as resulting from an eclipsed conformation of 1a. This interpretation is supported by the observation that the 25°C



1



2



3

Table I. Photochemical and Photophysical Data for Linked Anthracenes 1.

| | X | Y | ϕ_{12}^a | ϕ_{21}^a | ϕ_f^b | $\tau_f^b(\text{nsec})$ |
|-----|---------------------------------|----------------------------------|---------------|-------------------|----------------------|-------------------------|
| (a) | CH ₂ CH ₂ | H,H | 0.26 | 0.55 | 0.16 | 1.7 |
| (b) | CH ₂ CH ₂ | CH ₃ ,CH ₃ | 0.04 | --- | 0.14 | 2.0 |
| (c) | CH ₂ | H,H | 0.15 | 0.76 | 0.06 | 1.1 |
| (d) | CHOH | H,H | 0.29 | 0.81 | 0.02 | 0.3 |
| (e) | CHOH | H,OCH ₃ | 0.05 | --- | 0.02 | 0.3 |
| (f) | CH ₂ CH ₂ | CH ₂ CH ₂ | 0.36 | 0.60 ^c | < 0.001 ^d | (~0.01) ^e |


^aIsomerization quantum yield, benzene, 25°; 1 → 2, 366 nm, 2 → 1, 285 nm. ^bFluorescence yield and lifetime in benzene, 25° (single photon counting measurements except where noted). ^cRef. 1. ^dRef. 2. ^eFrom dimethylaniline quenching of photoisomerization (k_q assumed $\approx 10^{10} \text{ M}^{-1} \text{ sec}^{-1}$ as shown for 1a and 1d).

fluid solution fluorescence spectrum of 1a is abnormally broad and doesn't clearly show the characteristic vibrational bands noted in most anthracenes and in 1c-e.⁷ The ambient temperature spectrum is presumably that of averaged conformations. The observations concerning 1a are consistent with low temperature studies recently reported,⁸ in which two excimer-like conformations were assigned geometries with different degrees of interchromophore interaction.⁹

We conclude that linked anthracenes 1c-e are negligibly stabilized in the excited state due to interaction of the aromatic rings. On the other hand, a geometry which is nominally acceptable for excimerization is available to 1a. The striking result is that photoisomerization proceeds readily for these diverse systems (including 1f, where a high degree of chromophore interaction is revealed in absorption³), without

significant penalty for geometrical constraint in quantum efficiency or reaction rate.¹⁰ We conclude that a sandwich-like excimer state is not a necessary intermediate for anthracene photodimerization.

Diradicals. Linked anthracenes 1 have relatively short fluorescence lifetimes. The results in Table I show moreover that the rate of excited state decay is dependent on structure. With increasing anthracene proximity (note Table divisions), singlet lifetimes fall, but not in a way that can be rationalized by the yield of photoproduct. A mechanism, which accounts economically for the results, involves formation of an intermediate which partitions to reactant and photoisomer.



The ensuing kinetic relationships are $k_f = \Phi_f/\tau_f$, $k_r + k_d = 1 - \Phi_f/\tau_f$, and $P = \Phi_{12}/k_r\tau_f$, where k_f , k_r , and k_d are rate constants for fluorescence and reactive¹² and unreactive radiationless decay, respectively, and where P is a partition factor. If k_d is $< 10^8 \text{ sec}^{-1}$ (normally associated with intersystem crossing and triplet decay),^{4d,11,13} then reactive decay (k_r) is lifetime limiting and intermediate partitioning largely controls quantum efficiency.

The variety of geometrical constraints offered by the linked anthracenes (which significantly affects excited state decay rate but not quantum yield¹⁴), provides a clue to the nature of a sufficient intermediate for photoisomerization. Differences in the arrangement of anthracene moieties largely disappear with the formation of a 9,9' bond. Identification of intermediate I as diradical 3 would provide a unifying mechanism consistent with (1) singlet lifetime dependence on X , (2) lifetime independence on Y , and (3) quantum wastage. In short, the strict requirement for high

Table II. Kinetic Parameters for Photoisomerization of λ .^a

| | k_f | $k_r + k_d$ | p^b |
|-------------|-------|-------------|-------|
| λ_a | 0.94 | 4.9 | 0.31 |
| λ_b | 0.70 | 4.3 | 0.05 |
| λ_c | 0.55 | 8.6 | 0.16 |
| λ_d | 0.67 | 33 | 0.29 |
| λ_e | 0.67 | 33 | 0.05 |
| λ_f | < 1.0 | ~ 1000 | 0.36 |

^aRate constants $\times 10^{-8} \text{ sec}^{-1}$. ^bCalculated assuming $k_r \gg k_d$.

reactivity does not include a sandwich arrangement of aromatic rings or even a particular inter-ring dihedral angle, but more a proximity of two meso positions for sigma-bond formation. It follows that quantum efficiency for internal cycloaddition will depend largely on the nature of Y, and the effects observed in λ_b , λ_e , and λ_f are not surprising.

The proposal of diradical as sufficient intermediate in anthracene cycloaddition is supported by current theories concerning reactive radiationless decay.¹⁵ Intermediate λ is presumably the "pericyclic minimum" available to S_1 as the result of attempted crossing of ground and doubly excited states for the system. The theory permits that diradical λ be a common intermediate for $\lambda \rightarrow \lambda$ and $\lambda \rightarrow \lambda$, in which case $\phi_f + \phi_{12} + \phi_{21} = 1$ ¹ (excluding unreactive decay of λ). Within experimental error, our results are consistent with this scheme.¹⁶

That the geometrical requirements for excimerization¹⁷ exceed those for photo-reaction of the linked anthracenes suggests a subordinate role for excimers in

photodimerization but does not exclude their intermediacy in the general case. The data speak more to the nature of reactive decay of proximal anthracenes, whatever the degree of overall excited state stabilization of aromatic moieties due to excitation resonance^{4d} or other factors. The issues are: what is the minimum arrangement of anthracenes necessary for rapid photoreaction and, given that a number of geometries may be acceptable, is photodimer formation a one- or two-step process? The observed structure-reactivity relationships for the linked anthracenes provide valuable insight for both of these questions.

Acknowledgements. This work was supported by the Office of Naval Research and the National Science Foundation (through a fellowship to W.R.B.). We thank Professor M. Z. Hoffman for use of a fluorescence spectrophotometer and Professor A. M. Halpern for the lifetime measurements and most helpful discussions.

References and Notes

- (1) (a) G. Kaupp, Ann., 844 (1974); (b) G. Kaupp, Angew. Chem. Intl. Ed., 11, 313 (1972).
- (2) We observe (benzene solutions, in agreement with Kaupp¹) neither the fluorescence nor the wavelength dependence in photoisomerization quantum yield reported by Shizuka, et al.³ for THF solutions of 1f.
- (3) H. Shizuka, Y. Ishii, M. Hoshina, and T. Morita, J. Phys. Chem., 80, 30 (1976).
- (4) (a) M. D. Cohen, A. Ludmer, and V. Yakhot, Chem. Phys. Letters, 38, 3908 (1976); (b) A. Castellan, R. Lapouvade, and H. Bouas-Lourent, Bull. Soc. Chim. France, 210 (1976); (c) E. A. Chandross and J. Ferguson, J. Chem. Phys., 45, 3564 (1966); (d) J. Birks, "Photophysics of Aromatic Molecules," Wiley-Interscience, NY (1970) ch. 7; (e) J. Ferguson and A. W.-H. Mau, Mol. Phys., 27, 377 (1974); (f) J. Ferguson and S. E. H. Miller, Chem. Phys. Lett., 36, 635 (1975); and (g) B. Stevens, Adv. in Photochem., 8, 161 (1971). and (h) N.C. Yang, D.M. Shold, and B. Kim, J. Am. Chem. Soc., 98 6587 (1976).
- (5) (a) E.A. Chandross, J. Ferguson, and E.G. McRae, J. Chem. Phys. 45, 3546 (1966); (b) E.A. Chandross and J. Ferguson, J. Chem. Phys., 45, 3554 (1966).

- (6) J. Ferguson, A. W.-H. Mau, and J. M. Morris, Aust. J. Chem., **26**, 91, 102 (1973).
- (7) Room temperature solution absorption spectra for λ_{a-e} are nearly identical to that of 9-methylanthracene (or composite with 9-methoxy-10-methylanthracene) with slight alterations in vibronic intensities.
- (8) (a) J. Ferguson, M. Morita, and M. Puza, Chem. Phys. Lett., **42**, 288 (1976); (b) T. Hayashi, N. Mataga, Y. Sakata, S. Misumi, M. Morita, and J. Tanaka, J. Amer. Chem. Soc., **98**, 5910 (1976).
- (9) In addition to the structured "P band"^{8a} we have also observed the broad, structureless but very weak emission which occurs at longer wavelengths noted by Ferguson^{8a} and Mataga.^{8b}
- (10) Compare, for example, phenomenological rate constants for photoisomerization of λ_{c-e} (ϕ_{12}/τ_f , $0.1 - 1.0 \times 10^9 \text{ sec}^{-1}$) with pseudo first order rate constants for photodimerization of anthracene and 9-substituted anthracenes,¹¹ $0.4 - 1.0 \times 10^9 \text{ sec}^{-1}$ at 1 M, a concentration well above the level of complete concentration quenching of anthracene fluorescence (i.e., where virtually all excited state deactivation is bimolecular).
- (11) R. S. Shon, D. O. Cowan and W. W. Schmiegell, J. Phys. Chem., **79**, 2087 (1975).
- (12) For $\lambda_{a,b}$ k_r is a composite of conformer equilibrium constant and the rate constant for internal cycloaddition.
- (13) Sensitization and quenching experiments confirm that the involvement of triplets for λ is minimal.
- (14) The trend in fluorescence quantum yield and lifetime continues for a three atom linked system, 9,9'-dianthrylcarbonate ($\phi_{12} = 0.02$, $\tau_f = 5.3 \text{ nsec}$, $\phi_f = 0.30$). In this case, however, the yield of photoisomer is depleted significantly by competitive fluorescence and perhaps intersystem crossing. Presumably, diradical formation is slow due to an unfavorable entropy factor; i.e., in this case anthracene-anthracene orientation is important but more the availability than type of interchromophore arrangement. Excimer-type fluorescence was not observed for the carbonate.
- (15) (a) J. Michl, Top. in Current Chem., **46**, 1 (1974); (b) W. G. Dauben, L. Salem, and N. J. Turro, Accounts Chem. Research, **8**, 41 (1975).
- (16) Other features of the reaction coordinate include data for the thermal back reaction for selected systems: λ_a , $E_a = 32.8$, $\Delta H = -14.6$; λ_c , $E_a = 32.7$, $\Delta H = -17.3$; λ_f , $E_a = 22.3$, $\Delta H = -8.2$ (all in kcal/mol). The reaction enthalpies were determined by differential scanning calorimetry for isomerization in the solid state.

(17) Chandross and Dempster¹⁸ have shown that geometrical requirements are quite strict for excimerization of dinaphthylpropanes.

(18) E. A. Chandross and C. J. Dempster, J. Amer. Chem. Soc., 92, 3586 (1970).

William R. Bergmark*

Department of Chemistry

Ithaca College

Ithaca, New York 14850

Guilford Jones, II*

Department of Chemistry

Boston University

Boston, Massachusetts 02215